

had the lowest post-op outcome scores of all groups. This highlights the possibility that nutritional balance for patients is more than their BMI.

Arthroscopic Acetabular Labral Repair in Patients over the Age of 60: A Matched Case-Control Study

SS-33

May 19, 2017, 1:50 PM

J W THOMAS BYRD, M.D., PRESENTING AUTHOR
KAY JONES, M.S.N., R.N.



Introduction: The purpose of this study is to report the results of labral restoration among patients over age 60 compared to a cohort of younger adults.

Methods: 23 consecutive patients over age 60 undergoing labral repair with minimum one-year follow-up were compared (modified Harris hip Score) to a contemporaneous group of 23 patients age 18-55 matched for gender, degree of chondral damage, and associated FAI or dysplasia.

Results: Follow-up averaged 17.5 months (12 - 24 months). The study group consisted of 13 males and 10 females, average age 63 years (61 - 71); with 21 FAI and one dysplasia. 20 had acetabular articular damage (2 grade IV, 12 grade III, 5 grade II, 1 grade I) and 7 had femoral changes (1 grade IV, 6 grade 3). The exactly matched control group averaged 36 years (20 - 54). Study group improvement averaged 21 points (-19 - 46 points) with 20 (87%) improved. The average improvement in the control group was 19 points (-9 - 34) with 20 (87%) improved. There was no statistically significant difference between the two groups and the amount of improvement with statistically ($p < 0.01$) and clinically (> 5 points) significant in both. Two study group patients underwent THA at average 10 months with one control group THA at 11 months. All three converted to THA had combined grade IV acetabular and grade III femoral damage. There were no repeat arthroscopies and no complications in either group.

Conclusion: Patients over age 60 can benefit from arthroscopic labral repair with improved outcomes, modest rates of conversion to THA, and small risk of complication. Results are comparable to younger adults. Combined bipolar grade IV and grade III articular damage is a harbinger of conversion to THA, regardless of age.

Correlation of PROMIS CAT with Validated Hip Outcome Scores in Patients Undergoing Hip Arthroscopy

SS-34

May 19, 2017, 1:55 PM

SEAN CHILDS, M.D., PRESENTING AUTHOR
COLIN CANHAM, M.D.
RAYMOND KENNEY, M.D.
DIANA SILAS, D.O.
KELLY ADLER, M.ED., A.T.C.
BRIAN GIORDANO, M.D.



Introduction: The Patient Reported Outcomes Measurement Information System (PROMIS) tool was developed

by the National Institute of Health to provide efficient, precise and valid patient-reported outcome data. The purpose of this study was to determine whether the PROMIS Computer Adaptive Testing (CAT) tool demonstrates validity against commonly used legacy PRO measures among a population of patients undergoing arthroscopic hip surgery.

Methods: Eligible patients undergoing elective arthroscopic hip surgery completed a series of outcome questionnaires including the visual analog pain scale (VAS), a hip outcome score (HOS; ADL and Sports Subscales), modified harris hip (mHHS), and non-arthritic hip score (NAHS) as well as the PROMIS CAT for pain, physical function and emotional distress (depression). Patients completed all questionnaires at their preoperative evaluation and at regular postoperative intervals.

Results: A total of 77 patients elected to be included in this study, 28 (36.8%) were male and 49 (64.5%) were female. With regards to questionnaire completion, data from all patient reported outcome measures were gathered from 77 patients at their preoperative clinical visit, 71 during their first postoperative follow up, 69 at 6 weeks and 58 at 3 months postoperatively. Pearson correlations between PROMIS Physical Function T scores and the HOS ADL, HOS Sports, NAHS and mHHS were found to be 0.858, 0.799, 0.773 and 0.830 respectively. With regards to pain, the Pearson correlation between the PROMIS Pain T score and the VAS for pain was found to be 0.599.

Conclusion: The PROMIS tool was found to correlate well with current standards for patient reported outcome measures for individuals undergoing arthroscopic hip surgery. PROMIS Physical Function demonstrated high correlation with validated PRO. PROMIS Pain had moderate correlation with VAS Pain scores. With this information, the PROMIS tool can be used as a highly efficient and generalizable tool to collect pre- and post-operative data during clinical patient encounters.

The Relationship Between Arthroscopically Defined Acetabular Cartilage Defects and Preoperative dGEMRIC Indices: Refining the use of dGEMRIC

SS-35

May 19, 2017, 2:40 PM

ITAY PERETS, M.D., PRESENTING AUTHOR
EDWIN CHAHARBAKHSHI, B.S.
DAVID HARTIGAN, M.D.
BRIAN MU, B.A.
MARY CLOSE, B.S.
BENJAMIN DOMB, M.D.



Introduction: dGEMRIC is an advanced imaging technique that may detect early chondral damage. Although dGEMRIC may be useful in diagnosing early arthritis, there is a paucity of literature correlating local chondral damage with dGEMRIC indices. The purpose of this study was to report associations between dGEMRIC indices and intraoperatively defined acetabular cartilage damage in non-arthritic hips with femoro-acetabular impingement and/or labral tears, and to evaluate a new dGEMRIC index